

Order No. 704
K.H.L. King



No. 704

'Copymate' RECORD BOOK

Manufactured by the
Printing

50 leaves in Duplicate

6-12-1952

NUMBERED

NO CARBON PAPER REQUIRED

1871
1872

Journal of the ...

...

...

...

...

...

...

...

...

...

...

...

...

...

and the
the

the

the

the
the
the

the

the

the

the
the
the

the

the

the

the

the

the

Muntingia calabura F. Muell.

scattered patches of a lobate to 1-2ft (spec.)

Monstria mainly on this, ^{off a slender-leaved} ^{or a prostrate *Podocarpus*}

Cecropia 18 in to 3ft, individuals of ~~*Cecropia*~~

a pair of other shrubs, below end of road patches of very aromatic *Helichrysum hookeri* (Sond.) Bence & shrubby *Helichrysum* (?) (spec.) 1-3ft in bloom

(a less frequent host of *Monstria*), *Poa*

australis scattered everywhere & often in

large open areas, various forbs, including only one plant of *Cleome* spec.

below the end of road were

patches of *E. pauciflora* to 10ft & a seepage channel. Much the same shrubs as before

but growing to 3-4ft, also another

Phorbium phytolifolium F. Muell.

aromatic one (spec.) (no *Monstria*), the

seepage channel with epacrids, *Richea*,

Leptocarpum sp., patches of sedges &

hygrophilous forbs, patches of *Sphagnum*

Chaulioidium altatum not uncommon, ^{including nymphs.} One or

two *Monstria* were found amongst a mixture

of the *Cecropia* & pumy-fruited legume well

away from either of the favoured plants.

Return.



100 ft. to 150 ft. ...
 ...
 ... to 100 ft. ...

... to 15 ft. ...

... to 2 ft. ...

... 2 ft. ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

...
 ...

ociput, from antennal scrobes below
 eyes & along ventral margin of pronotal
 lobe, & ventral margin of femur III.

Genicular area of femur III bare. Abdomen
 cephalad under & adjacent to wing ^{usually}
 smooth & scarlet. Antennae black.]

saw one male settle after disturbance

& vibrate the antennae while raising

the tegmina ^{dorsolaterally} to expose the wings. Very

few females seen (2) - presumably flying

to eggs in shrubs. No *Parasitica* seen below

the herb field, although a fair amount

of *Parabebe* was present in places in

the *E. delegatensis* forest.]

8/2

7.3 Mansfield's depart.

9.3 Jamieson. Return.

11.0 Turned left on Eildon road.

12.0 Cummins' Gap. Stop slopes in
Stenobothrus vittatus, *Parabebe*, *Parasitica*,
P. obliqua, *E. n. acrostychna*, *E. n.*

many others to left.

1. The site simple, for the purpose of
 the *Trichostema* *Trichostema* *Trichostema*
Trichostema & *E. binnendijkii* *Trichostema*

2. The *Trichostema* *Trichostema* *Trichostema*
 of, also, *Trichostema* *Trichostema* *Trichostema*
 lead off to

3. *Trichostema* *Trichostema* *Trichostema*

4. *Trichostema* *Trichostema* *Trichostema*

5. *Trichostema* *Trichostema* *Trichostema*

6. *Trichostema* *Trichostema* *Trichostema*

7. *Trichostema* *Trichostema* *Trichostema*
Trichostema

8. *Trichostema* *Trichostema* *Trichostema*
Trichostema

9. *Trichostema* *Trichostema* *Trichostema*
Trichostema

10. *Trichostema* *Trichostema* *Trichostema*
Trichostema

11. *Trichostema* *Trichostema* *Trichostema*
Trichostema

12. *Trichostema* *Trichostema* *Trichostema*
Trichostema

13. *Trichostema* *Trichostema* *Trichostema*
Trichostema

14. *Trichostema* *Trichostema* *Trichostema*
Trichostema

15. *Trichostema* *Trichostema* *Trichostema*
Trichostema

24. Walden, Idaho.

25. Thompson, Tenn & left.

26. Tugarty, Tenn & left.

27. Garrett, Tenn & left.

28. Morgan, W. Va.

29.

30. Marionville, Mo.

31. Tenn & left.

4433 mi.

32. Summit, Lake Mountain of
Altoona, Pa. to the base of the mountain
to the base of the mountain
to the base of the mountain
to the base of the mountain

to a fairly good degree.

to a fairly good degree.

to a fairly good degree.

to a fairly good degree.

33. Tenn & left to the mountain

to a fairly good degree.

to a fairly good degree.

to a fairly good degree.

The first patch of open track was
 the surface scattered about some of
 the stones after a short parallel gap
 that similar green after a nymphs
 ground in the vicinity of the
 (some faintly leafless). The
 population was sparse they were
 flat spots beyond the peak for the
 females very green, the parts mostly
 on the shores or on the ground
 the sign of *Monistria canina* -
 present, showing no evident
 any one species of shrub. *Rossia*
 all females seen nymphs
 the. Females in places, not forming
 abundant, also *Staphylinus*
 the following table sp. on some flat
 General account of *Monistria*
canina was not found below the
 low on hills, although *canina*
 spent several stops were
 essential at a certain time

Mimyacris nana

452.6 Total River.

Prostanthera sp. *Prostanthera* *mitis*, *Prostanthera*
 452.6 Total River. *Prostanthera* *mitis* *Prostanthera*
Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*
Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

452.6 Total River. *Prostanthera* *mitis* *Prostanthera* *mitis* *Prostanthera*
Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*
Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*
Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

452.6 Total River depart.

[Reflections on Wilson's Prostanthera]

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

Prostanthera *mitis* *Prostanthera* *mitis* *Prostanthera*

12/2

65 - 2nd floor.66 - 2nd floor.67 - 2nd floor.68 - 2nd floor.69 - Entered elevator to 2nd floor.70 - Took back to 2nd floor.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

ulnaris

Minyasius rana

23.5. Slight depression with the following
 grasses & herbs - *usitatus*
 shrubs to 2-3ft interspersed with
 grass & herbs - *Koeleria* *laevis*
 shrubs.

23.6. Slight depression. E. part. *Panicum*
virgatum *Panicum* *capillare* *Panicum*
 with several spp. of shrubs to 3ft.

abundant silver-leaved *Clusia*.

26.4. end of Barro Colorado. *Heisteria*.

27.2. steep slope with E. part. *Panicum*
Momystica *concinna*, *Panicum* *capillare*
 several other species, including *Heisteria*.

with *Panicum* *capillare* *concinna* to *Momystica*.

roughly 200m.

28.9. *Panicum* *capillare* *concinna* - *Panicum* *capillare*